

CROSSBREEDING PARAMETERS FOR GROWTH PERFORMANCE OF BROILER RABBITS*

L. Zavadilová¹, K. Mach², I. Majzlík²

¹Research Institute of Animal Production, Prague-Uhřetěves, Czech Republic

²Czech University of Agriculture, Faculty of Agrobiological Sciences, Prague, Czech Republic

The growth of broiler rabbits: two parental lines 59 (paternal) and 19 (maternal) and three groups of crossbreds F1 (59 x 19), F2, 59 x F1 was analysed in six week fattening period (42 d to 84 d of age). The analysed traits were live weight (BW), the average daily gain (ADG), average daily feed consumption (ADF) and feed conversion (FC) (weekly). The estimated least squares means were used as input data for estimating the crossbreeding effects. The model of Dickerson (1969, 1973) was used in all calculations. Three kinds of crossbreeding effects were included: additive effect, heterosis and recombination loss. By the end of the fattening period, the heaviest crossbreds F1 attained 2802.5 g and paternal line 59 2935.8 g whereas the lowest BW was noticed for crossbreds F2 (2606 g). For BW direct additive effects of line 59 were from 1.5 g to 321.4 g during the whole fattening period. Heterosis was negative and non-significant (-10.37 g to -112.14 g). Recombination loss occurred negative, high and significant (-14.72 g to -493.51 g). Significant additive effect for ADG was found only in the first week of test (20.34 g), significant heterosis occurred in the third week of experiment (3.29 g). Recombination loss was in the range from -23.86 g to 16.92 g. Additive effects for ADF were non-significant (-23.03 g to 51.11 g) the heterosis was significant and mostly positive (37.03 g to 42.39 g). The recombination loss ranged between -74.10 g and 55.70 g. The estimates of crossbreeding effects for FC were small and non-significant.

rabbit; growth; feed conversion; crossbreeding; maternal line; paternal line

INTRODUCTION

The broiler rabbit is generally a two-way or a four-way hybrid. A performance of these hybrids is determined by several crossbreeding effects, especially by heterosis. In addition, crossbreeding with specialized breeds or strains make use of the complementarity between dam lines, which have a good reproductive efficiency, and sire lines, which possess a good meat production efficiency. Knowledge of a magnitude and a direction of crossbreeding effects are important for crossbreeding scheme decision.

The testing of broiler rabbits has been conducted by the Department of Genetics and Animal Production of the Czech University of Agriculture in Prague since 1992. The testing is concentrated on growth, feed consumption, feed conversion and carcass quality (Mach, 1992; Mach, Majzlík, 1993; Mach et al., 1997a, b; Dědková et al., 1999, 2002).

The objective of the present study was the estimation of crossbreeding effects for growth performance of two parental lines and their crossbreds during the fattening period from 42 days to 84 days of age.

MATERIAL AND METHOD

The growth of broiler rabbits: two parental lines 59 (paternal) and 19 (maternal) and three groups of cross-

breds F1 (59 x 19), F2, 59 x F1 was analysed. The test was carried out at the Department of Genetics and General Animal Science of Czech University of Agriculture in Prague. The six week long fattening period started at 42 days and finished at 84 days of age of the rabbits. The live weight (BW) and individual feed consumption was weekly measured until the end of the experiment. The average daily gain (ADG), average daily feed consumption (ADF) and feed conversion (FC) was calculated from data for every experimental week.

The analysis was carried out by the ordinary least squares method in the procedure GLM of statistical software SAS (SAS, 1988). A linear model included the factors of the rabbit group and factor of breeder. The estimated least squares means were used as input data for the program package CBE4, version 4.0 (Wolf, 1995) that was used for estimating the crossbreeding effects. The model of Dickerson (1969, 1973) was used in all calculations. Three kinds of crossbreeding effects were included: additive, heterotic and recombination effects. The additive effects for maternal line 19 were set to zero. The additive effects for paternal line 59 were therefore expressed as deviations from maternal line.

The coefficients for these effects were expressed as given in Wolf et al. (1995). The full model for crossbreeding effects has the following form then:

$$CBE = \alpha_{59} \alpha_{59} + \delta_{19,59} h_{19,59} + (4\alpha_{19} \alpha_{59} - \delta_{19,59}) r_{19,59}$$

* The research was supported by the Ministry of Agriculture of the Czech Republic (Project MZE 0002701401) and Ministry of Education, Youth and Sports of the Czech Republic MSM 6046070901.

where: a – additive effect of line 59
 h – heterosis for combination 19 x 59
 r – recombination loss for breed combination 19 x 59

The α 's and δ 's are the coefficients, they are specific for each animal.

The estimation was carried out by weighted least squares using the reciprocals of the variances of the genetic group means as weights.

RESULTS AND DISCUSSION

By the end of the fattening period, the heaviest crossbreds F1 attained 2802.5 g, paternal line 59 2935.8 g, whereas the lowest BW was noticed for crossbreds F2 (2606 g) (Fig. 1).

The ADG (Fig. 2) dropped off within weeks of the fattening period for all the groups of rabbits. The highest ADG was found for paternal line 59 (51.79 g), for cross-

breeds F1 (45.16 g) and 59xF1 (44.07g) in the first week of the test. For crossbreds F2 the ADG varied between weeks of fattening period.

Feed consumption (ADF) was proportional to the ADG of rabbits. The highest ADF (Fig. 3) was found for line 59 (141.79 g) and crossbreds F1 (137.19 g) and 59 x F1 (143.44 g) in the first week of fattening. The values of ADF fluctuated for crossbreds F2.

No marked differences were found for FC among all groups (2.90 to 5.98) (Fig. 4). An increase of FC was observed with the age of rabbits, especially in the last week of fattening period (max. 12.52). We cannot confirm findings of Brun and Ouhayoun (1989) that feed efficiency is reduced by crossbreeding.

The estimates for crossbreeding effects are presented in Table 1. Positive estimates of additive effect on behalf of line 59 were found for BW (31.5 g to 321.4 g) during the whole fattening period. It is in agreement with Dědková et al (2002). Heterosis for BW was negative




Fig. 1. Body weight of rabbits for the fattening period




Fig. 2. Average daily gain of rabbits for the fattening period




Fig. 3. Average daily feed consumption of rabbits for the fattening period




Fig. 4. Feed conversion of rabbits for the fattening period

and non-significant (-10.37 g to -112.14 g). In contrary, Mach, Majzlík (1993) and Brun, Ouhayoun (1989) found positive heterosis for BW at the end of fattening period (91 days of age, 79 days of age, respectively). Recombination loss occurred negative, significant and high (14,72 g to -493,51 g).

The crossbreeding effects for the rest of the analysed traits did not show obvious trends. The ADG was found to have significant additive effect only in the first week of test (20.34 g), significant heterosis occurred in the third week of experiment (3.29 g). The positive heterosis for daily gain was reported by Brun and Ouhayoun (1989) or Krogmeier and Dzapo (1991). Recombination loss was in the range from -23.86 g to 16,92 g. While the additive effects for ADF were non-significant (-23.03 g to 51.11 g) the heterosis turned out to be significant and mostly positive (37.03 g to 42.39 g). The es-

timates of recombination loss fluctuated between -74.10 g and 55.70 g. The estimates of crossbreeding effects for FC were small except for two of them which were non-significant.

We can conclude that the crossbreeding effects mostly influenced only body weight of the rabbits in this experiment. The positive additive effect increased the BW of rabbits proportionally to the part of the 59 line gene. On the contrary, the negative estimates of recombination loss restrained the BW of crossbreds especially in the F2.

Acknowledgements

We are grateful to J. Wolf for the program package CBE and for valuable discussions and to Barbora Hofmanová for technical assistance.

Table 1. Estimates of crossbreeding effects for analysed traits

Age (days)	Additive effect	SD		Heterosis	SD		Recombination loss	SD	
Body weight (g)									
42	31.49	64.446	ns	-10.37	39.132	ns	-14.72	86.553	ns
49	257.59	73.833	**	-46.77	46.214	ns	-326.80	86.830	**
56	292.23	85.463	**	-112.14	53.494	*	-472.13	100.507	**
63	304.15	90.967	**	-69.27	56.939	ns	-416.37	106.980	**
70	288.88	104.818	**	-36.88	65.608	ns	-493.51	123.269	**
77	303.46	106.585	**	-14.68	66.715	ns	-413.28	125.347	**
84	321.41	113.712	**	-24.65	71.175	ns	-482.15	133.728	**
Average daily gain (g)									
42-49	20.34	5.587	**	-2.38	3.393	ns	-23.86	7.503	**
49-56	4.85	5.587	ns	-1.04	3.393	ns	-11.49	7.503	ns
56-63	0.19	5.262	ns	6.48	3.292	*	8.57	6.213	ns
63-70	-1.04	5.014	ns	3.03	3.140	ns	-9.84	5.951	ns
70-77	1.02	6.530	ns	3.42	4.083	ns	16.92	7.759	*
77-84	1.75	5.623	ns	-1.24	3.511	ns	-7.06	6.742	ns
Average daily feed consumption (g)									
42-49	51.11	11.526	**	37.03	6.999	**	35.69	15.480	ns
49-56	13.83	15.215	ns	-21.26	9.523	*	-74.10	17.893	**
56-63	22.62	12.914	ns	36.66	8.084	**	30.28	15.188	ns
63-70	23.80	13.718	ns	40.44	8.587	**	-55.70	16.133	**
70-77	-6.28	14.148	ns	42.39	8.923	**	10.46	16.674	ns
77-84	-23.03	20.007	ns	-52.75	12.523	**	-97.62	23.529	ns
Feed conversion									
42-49	-0.78	0.804	ns	0.70	0.488	ns	2.33	1.080	*
49-56	0.66	1.482	ns	2.58	0.930	**	2.39	1.745	ns
56-63	0.44	1.306	ns	0.44	0.817	ns	1.12	1.543	ns
63-70	0.84	1.743	ns	1.69	1.092	ns	1.09	2.069	ns
70-77	-0.69	1.274	ns	0.92	0.796	ns	-0.42	1.513	ns
77-84	0.62	5.000	ns	-1.88	3.122	ns	-6.18	5.995	ns

SD – standard deviation, ns – non-significant, * $P < 0.05$, ** $P < 0.01$

REFERENCES

- BRUN, J. M. – OUHAYOUN, J.: Growth performance and carcass traits in three strains of rabbits and their two-way crosses. *An. Zootech.*, 38, 1989: 171–179.
- DĚDKOVÁ, L. – MACH, K. – MAJZLÍK, I.: Growth and feed conversion in broiler rabbits. *Scientia Agric. Bohem.*, 30, 1999: 315–323.
- DĚDKOVÁ, L. – MACH, K. – MAJZLÍK, I. – MOHSEN, A.: Analysis of growth and feed conversion in broiler rabbits by factorial crossing. *Czech J. Anim. Sci.*, 2002: 133–140.
- DICKERSON, G. E.: Experimental approaches in utilizing breed resources. *Anim. Breed. Abstr.*, 37, 1969: 191–202.
- DICKERSON, G. E.: Inbreeding and heterosis in animals. In: *Proc. of Animal Breeding and Genetics Symposium in Honor of Dr. Jay L. Lush. American Soc. Anim. Sci.*, 1973: 54–77.
- SAS: SAS-STAT. User's Guide, 6.03 Ed. SAS Institute, Inc., Cary, NC. 1988. 1028 pp.
- MACH, K.: Některé genetické a chovatelské aspekty masné užitkovosti králíků (Some genetic and breeding aspects of meat performance of rabbits). [Thesis for habilitation.] Prague, CUA 1992. 72 pp.
- MACH, K. – MAJZLÍK, I.: Uplatnění dialelního křížení při testaci masných plemen králíků (Application of diallel crossing in testing of meat breeds of rabbits). In: *Proc. of XVIth Days of Genetic of Livestock, Nitra, SR, 6–8 September 1993: 219–223.*
- MACH, K. – MAJZLÍK, I. – ŘÍČAŘ, L.: Testace výkrmnosti a jatečné hodnoty finálních hybridů brojlerového králíka (Testing of fattening and carcass value of final hybrids of broiler rabbits). In: *Sbor. přednášek IV. celostátního semináře "Nové směry v chovu brojlerových králíků" (Proc. of lectures of the IVth national seminar "New trends in broiler rabbit breeding")*, CUA, Prague, 19. 11. 1997a: 46–49.
- MACH, K. – MAJZLÍK, I. – KODEŠ, A. – RAIFOVÁ, I.: Rast, spotřeba krmiva a jatočná hodnota brojlerových

králíků (Growth, feed consumption and carcass value of rabbit broilers). In: Zbor. ref. z konferencie s medzinárodnou účasťou poriadanej pri príležitosti 50. výročia založenia VÚŽV v Nitre, 2. časť, 8.-9. 10. 1997b: 172-174.

KROGMEIER, D. – DZAPO, V.: Leistung Merkmale von Kaninchen der Rassen Weisse Neuseeländer, Helle Grosilber und deren reziproker Kreuzungen. 2. Mitteilung: Heterosissteigerungen in Mastleistungs-, Schlachtkörperqualitäts- und Fleischbeschaffenheitsmerkmalen. Arch. Geflügelk., 55, 1991: 162-169.

WOLF, J.: CBE – a general program for estimating crossbreeding parameters. In: 46th Annual Meeting of EAAP, 4-7 September, Prague, 1995.

WOLF, J. – DISTL, O. – HYÁNEK, J. – GROßHANS, T. – SEELAND, G.: Crossbreeding in farm animals. V. Analysis of crossbreeding plans with secondary crossbred generations. J. Anim. Breed. Genet., 112, 1995: 81-94.

Received for publication on March 13, 2005

Accepted for publication on May 5, 2005

ZAVADILOVÁ, L. – MACH, K. – MAJZLÍK, I. (Výzkumný ústav živočišné výroby, Praha-Uhřetěves; Česká zemědělská univerzita, Fakulta agrobiologie, potravinových a přírodních zdrojů, Praha, Česká republika):

Efekty hybridizace v růstu brojlerových králíků.

Scientia Agric. Bohem., 36, 2005: 87-91.

Byl vyhodnocen růst brojlerových králíků během šestitýdenního výkrmu (42 až 84 dní věku) u otcovské linie 59 a mateřské linie 19 a jejich kříženců F1 (59 x 19), F2 a 59 x F1. Pro živou hmotnost (BW), průměrný denní přírůstek (ADG), průměrnou denní spotřebu krmiva (ADF) a konverzi krmiva (FC) byly v každém týdnu výkrmu na základě průměrů nejmenších čtverců odhadnuty efekty křížení. Efekty křížení byly definovány podle modelu D i c k e r s o n a (1969, 1973). Odhadnut byl aditivní efekt, heterozní efekt a rekombinační ztráta. Na konci výkrmu dosáhli největší BW kříženci F1 (2802,5 g) a králíci otcovské linie 59 (2935,8 g), zatímco nejnižší BW byla nalezena u kříženců F2 (2606 g). Aditivní efekt linie 59 se pohyboval mezi 1,5 g až 321,4 g během celé doby výkrmu. U BW byl heterozní efekt negativní a statisticky nevýznamný (-10,37 g až -112,14 g). Rekombinační ztráta pro BW byla negativní a statisticky významná (-14,72 g až -493,51 g). Statisticky významný aditivní efekt pro ADG byl nalezen pouze v prvním týdnu testu (20,34 g). Podobně se statisticky významný heterozní efekt pro ADF objevil jen v třetím týdnu experimentu (3,29 g). Rekombinační ztráta se pohybovala v rozmezí od -23,86 g do 16,92 g. Aditivní efekty pro ADF byly statisticky nevýznamné (-23,03 g až 51,11 g), heterozní efekty naopak signifikantní a většinou kladné (37,03 g až 42,39 g). Rekombinační ztráta se pohybovala mezi -74,10 g a 55,70 g. Odhady efektů křížení pro FC byly malé a statisticky nevýznamné.

králík; růst; konverze krmiva; křížení; mateřská linie; otcovská linie

Contact Address:

Ing. Ludmila Zavadilová, Výzkumný ústav živočišné výroby, Praha-Uhřetěves, Přátelství 815, 104 01 Praha-Uhřetěves, Česká republika, e-mail: dedkova.ludmila@vuzv.cz
