

PHENOLIC COMPOUNDS IN WETLAND MACROPHYTES*

T. Dvořáková Březinová, J. Vymazal

Czech University of Life Sciences Prague, Faculty of Environmental Sciences, Prague, Czech Republic

Phenolics are the most widely distributed class of plant secondary metabolites and higher plants are able to produce several thousand different phenolic compounds. It has been reported that phenolics are more resistant to decomposition due to the negative impact on the microorganisms involved. Therefore, it may be hypothesized that the higher content of phenolics in plant tissue may cause slower decomposition rates and potentially higher carbon sequestration in the soil. The primary goal of this study was to evaluate and compare the content of phenolics in seven common macrophytes in the Czech Republic. Aboveground biomass was sampled in June 2016 at seven different localities and phenolics concentrations determined by spectrophotometric methods according to the Folin-Ciocalteu method ranged from 9.02 to 28.39 g kg⁻¹ dry biomass weight (DW). Several plants were also harvested in August, October and December in order to follow a seasonal pattern. According to the results it seems that surveyed macrophytes vary widely in phenolics concentrations in relation to sampled site, harvesting time, plant species, and plant parts as well.

phenolic compounds, wetland plant species, decomposition, carbon



doi: 10.2478/sab-2018-0001

Received for publication on July 24, 2017

Accepted for publication on September 14, 2017

INTRODUCTION

Phenolic compounds represent a large group of substances with diverse properties (Klaa et al., 2016). They are the most widely distributed class of plant secondary metabolites and they are very important for plant biochemistry and physiology. Phenolics are involved in many interactions of plants with the biotic and abiotic environment and plants can synthesize phenolic compounds as a reaction to changing environmental conditions (Hutzler et al., 1998). It has been reported that plants are able to produce several thousand different phenolic compounds (Lattanzio, 2013). Phenolics in plant play many important roles, which can be categorized into several groups. They can (a) be involved in growth and reproduction, (b) contribute to plant morphology and sensorial properties

(pigmentation, aroma, taste) (Bravo, 1998; Giada, 2013), and (c) provide passive and active resistance and protection against pathogens, predators or stress, and ultra violet radiation (Richardson et al., 1999; Vermerris, Nicholson, 2006a).

Structurally, these compounds have one or more hydroxyl group attached directly to the aromatic ring and the most basic phenolic compound is phenol. The terms polyphenols depicts compounds that have more than one phenolic hydroxyl group attached to one or more benzene rings (Vermerris, Nicholson, 2006b). A large number of these compounds can be divided into several classes and they can be classified in different ways – for example according to the number of constitutive carbon atoms or the structure of the basic skeleton (Murkovic, 2003). In general, phenolics range from simple phenols to highly polymerized

* Supported. by the Internal Grant Agency of the Czech University of Life Sciences Prague (CIGA), Project No. 20164207

Table 1. Surveyed macrophytes and sampling sites characteristics. 1-*Phragmites australis*, 2-*Phalaris arundinaca*, 3-*Typha latifolia*, 4-*Glyceria maxima*, 5-*Scirpus sylvaticus*, 6-*Carex nigra*, 7-*Juncus effusus*

Location	Type of wetland	Monitored plants
Březnice	wet meadow	1,2,3,4,5,6
Běloky	floodplain	1,2,3,5,7
Chmelná	floodplain	1,2,3,5,6,7
Horní Bradlo	stream bank	2,3,5,7
Louňovák (pond)	pond littoral zone	1,2,4,6
Moraveč	riparian wetland	2,5,7
Pařez (pond)	pond littoral zone	1,2,4,5,6

compounds like tannins or lignins (Bravo, 1998). In most cases, they do not occur as free compounds in plants, but they are bound to other molecules and are present as esters or glycosides (Vermerris, Nicholson, 2006b).

The levels of phenolics in plants differ among various species (Hoorens et al., 2003; Wang et al., 2015; Rejmankova, 2016; Harrison et al., 2017) and even among varieties of the same species (Bravo, 1998). The concentration of phenolics is partly influenced by genetic background and partly by environmental factors (Connor et al., 2002; Howard et al., 2003), especially the nutrients availability. Harrison et al. (2017) concluded that the variability was influenced by local environmental factors and the most important predictive factors for foliar phenolic compounds in some wetland species were sampling date, soil nutrients, and herbivory. Rejmankova (2016) showed that the concentration of phenolics in *Eleocharis cellulosa* and *Typha domingensis* were negatively correlated with increased growth due to increasing nutrient levels. She confirmed the protein competition model (PCM) proposed by Jones, Hartley (1999) which assumes that protein and phenolics synthesis compete for the common limiting resource phenylalanine and, therefore, protein and phenolics allocation are inversely correlated. Under suitable nutrient conditions, most of the phenylalanine is used for the formation of proteins, while during the nutrient limiting conditions phenylalanine is more used for the formation of phenolics (Rejmankova, 2016). The concentration of phenolics can differ even in different parts of the same plant. For example, formation of flavonol and flavone glycoside in common vegetables depends on light, so these compounds are mostly concentrated in outers organs like leaves than in any other part of the same plant (Herrmann, 1988).

Polyphenols have been recognized as regulators of soil processes and it has been suggested that they inhibit nitrification, as well as plant litter decomposition and nutrient recycling (Hornner et al., 1988; Xu et al., 2013). Phenolics, such as lignins, unlike proteins and carbohydrates, are more resistant to decomposition due to the negative impact on the microorganisms involved

in the decomposition of plants and, therefore, phenols can accumulate temporarily in soil. In the environment, soluble polyphenols face four different fates. They can be: (a) degraded and mineralized as a carbon source by heterotrophic microorganisms, (b) transformed into insoluble and recalcitrant humic substances by polymerization and condensation reactions (with the contribution of soil organisms), (c) adsorbed to clay minerals or chelated with Al or Fe ions, or (d) leached by percolating water, and leave the ecosystem as part of dissolved organic carbon (DOC).

The effects of polyphenols on soil microorganisms were reviewed by many researchers, for example by Kuiters (1990), Hattenschwiler, Vitousek (2000) or Wang et al. (2015). According to these studies the most resistant compounds to decompose in plants are phenolics and, therefore, it may be hypothesized that higher content of phenolics in plant tissue leads to slower decomposition rates. It can also be expected that concentrations of phenolics may play an important role in carbon sequestration in wetlands, and consequently contribute to mitigation of global warming effects.

Decomposition is very important in wetlands, because wetlands have been proved to be one of the major carbon sinks, mainly due to the incomplete decomposition of plant material. In wetlands, the term decomposition is mostly confined to the breakdown and subsequent decay of dominant macrophytes, which leads to the production of detritus (Boyd, 1970). Most annual plant production in wetlands is not consumed by herbivores due to the poor digestibility, thus it decomposes on the wetland surface and becomes a part of a particulate carbon pool (Teal, 1962; Gallagher, 1978; Polunin, 1982).

Despite this significant fact, studies focused on phenolics concentrations in wetland macrophytes are very limited. Studies by Rejmankova (2016), Wang et al. (2015), and Harrison et al. (2017) focused on wetland macrophyte species in salt, brackish, and freshwater wetlands in North and Central America, whereas information about this topic in European climate conditions is limited to the information from the Netherlands and northern Sweden (Hoorens et al., 2003; Dorrepall, 2005; Aerts et al., 2006). Hence, the primary goal of this study was to evaluate and compare the content of phenolics in seven common wetland macrophytes in the Czech Republic and thus extend the knowledge about phenolics content in-macrophytes.

MATERIAL AND METHODS

Study sites and plant sampling

Aboveground biomass of seven common wetland macrophytes was sampled at seven different localities

across the Czech Republic (Table 1). The macrophytes involved in the study were *Phragmites australis* (Cav.) Trin. ex Steud., *Phalaris arundinacea* L., *Typha latifolia* L., *Glyceria maxima* (Hartman) Holmberg, *Scirpus sylvaticus* L., *Carex nigra* (L.) Reichard, and *Juncus effusus* L. In *P. australis* and *P. arundinacea*, leaves and stems were analyzed separately. In *T. latifolia*, only leaves were analyzed as flower-stems did not occur at all locations. In order to eliminate varying concentrations in different parts of the leaves and stems, the whole stems and leaves were analyzed. The study was carried out in 2016 and all surveyed plants were harvested in the last week of June. Several plants were also harvested in August, October, and December in order to follow a seasonal pattern. All samples consisted of three plants and all samples were taken in three replicates.

Analytical methods

Harvested biomass was separated into stems and leaves (in the case of *P. australis* and *P. arundinacea*), dried to a constant weight at 60°C, and ground in the cutting mill Pulverisette 15 (Fritsch, Idar-Oberstein, Germany) using the 0.5 mm mesh size screen. The ground plant material (approximately 0.1 g) was extracted in 5 ml of 70% acetone for 1 h at 4°C. Total content of phenolics was determined by spectrophotometric methods according to the Folin-Cioacalteau method (Barlocher, Graca, 2005). Standards solutions ranging from 0 to 250 mg l⁻¹ were prepared from tannic acid (Sigma Aldrich, St. Louis, USA) dissolved in 70% acetone. Standards solutions and spine

extracts were transferred to the test tubes, solution of 2% Na₂CO₃ in 0.1 M NaOH and Folin-Cioacalteau reagent (Sigma Aldrich) diluted 1 : 2 with deionized water were added. Absorbance at 760 nm was measured after 120 min using Cary UV-Vis 60 spectrophotometer (Agilent Technologies, Santa Clara, USA). The phenolics concentration was expressed in g per kg dry biomass weight (DW).

The statistical analysis of variance (ANOVA) followed by post-hoc Tukey's HSD test was used to evaluate differences between phenolics contents of various plants. The significance level was set at $P < 0.05$.

RESULTS

Total phenolics in macrophytes

The average concentrations of phenolics in above-ground biomass of all monitored plants are shown in Fig. 1. The concentrations of total phenolics varied among surveyed species and in various plant parts as well. The lowest concentrations were found in stems of *P. arundinacea* (9.02 ± 0.14 g kg⁻¹ DW) followed by stems of *P. australis* (10.47 ± 0.78 g kg⁻¹ DW). The highest concentrations were observed for *Scirpus sylvaticus* (27.74 ± 0.96 g kg⁻¹ DW) and *Carex nigra* (28.39 ± 0.54 g kg⁻¹ DW).

Seasonal pattern of total phenolics concentrations

The results revealed several different seasonal patterns of phenolics concentrations in aboveground




Fig. 1. Average concentrations (\pm SD) of total phenolic compounds in aboveground biomass of seven macrophytes: *Phalaris arundinacea* ($n = 7$), *Phragmites australis* ($n = 5$), *Glyceria maxima* ($n = 3$), *Typha latifolia* ($n = 4$), *Juncus effusus* ($n = 4$), *Scirpus sylvaticus* ($n = 6$), *Carex nigra* ($n = 4$) harvested in June 2016

DW = dry biomass weight; ^{a-c}different letters indicate a significant difference at $\alpha = 0.05$ between the means across species




Fig. 2. Concentrations of total phenolic compounds in aboveground biomass of *Scirpus sylvaticus* sampled during the period June–December 2016; data presented are means \pm SD DW = dry biomass weight

^{a,b}different letters indicate a significant difference at $\alpha = 0.05$ between the means

biomass. The concentrations of total phenolics in aboveground biomass of *Scirpus sylvaticus* sampled during the period June–December 2016 are shown in Fig. 2. The concentration decreased from the highest value in June ($27.84 \pm 1.53 \text{ g kg}^{-1}$) to the lowest one recorded in October ($7.88 \pm 0.733 \text{ g kg}^{-1}$) and remained low until December. Similar behaviour was observed in leaves of *Phalaris arundinacea* (Fig. 3). The average concentration was the highest and steady in June ($19.36 \pm 0.84 \text{ g kg}^{-1}$) and during the summer months, then the concentration decreased until December, where the lowest value ($7.03 \pm 0.30 \text{ g kg}^{-1}$) was observed. On the other hand, the seasonal pattern of phenolics concentration in stems of the same plant exhibited a different course (Fig. 3). The average concentrations also decreased from June to December, but the decline was very slow, from 9.97 ± 0.06 to $6.07 \pm 0.57 \text{ g kg}^{-1}$.

A different seasonal pattern was observed for total phenolics concentration in *Typha latifolia* (Fig. 4). The highest value was recorded in August ($18.12 \pm 2.85 \text{ g kg}^{-1}$) while the lowest value was observed in December ($14.52 \pm 2.99 \text{ g kg}^{-1}$). However, the differences between the average concentrations of each month are not significant.

Concentration of total phenolics in macrophytes at different localities

Phenolics concentrations in leaves and stems of *Phragmites australis* sampled in June at five different natural wetlands are shown in Fig. 5A. The highest average concentration in leaves ($21.53 \pm 1.16 \text{ g kg}^{-1}$) was recorded in Chmelná. The lowest value was observed

in Březnice ($16.38 \pm 1.26 \text{ g kg}^{-1}$), but statistically, no significant difference between means was proved. In the case of stems, concentrations varied significantly between the highest one in Louňovák ($12.44 \pm 0.31 \text{ g kg}^{-1}$) and the lowest one in Březnice ($8.92 \pm 0.64 \text{ g kg}^{-1}$). Differences in phenolics concentrations between the highest and the lowest values for leaves and stems amounted to 25 and 28%, respectively. A similar pattern was found for *P. arundinacea* (Fig. 5B), where no significant differences were observed between the highest (22.50 g kg^{-1} in Moraveč) and the lowest (15.26 g kg^{-1} in Horní Bradlo) values in the leaves. Phenolics concentrations in stems varied between only $9.97 \pm 0.06 \text{ g kg}^{-1}$ in Březnice and $7.99 \pm 0.26 \text{ g kg}^{-1}$ in Chmelná but the significant differences were recorded.

Carex nigra (Fig. 5C) exhibited very similar phenolics concentrations at three locations – 28.92 ± 0.39 , 28.50 ± 0.86 , and $26.66 \pm 0.33 \text{ g kg}^{-1}$ at Březnice, Pařez and Louňovák, respectively. A significantly lower value was observed in Chmelná, where the concentration reached only $10.86 \pm 0.92 \text{ g kg}^{-1}$. This value was by 62% lower as compared to that at Březnice.

A similar pattern was observed for *Typha latifolia* (Fig. 5D) – the highest concentrations were observed in Běloky ($23.52 \pm 6.32 \text{ g kg}^{-1}$) and Horní Bradlo ($23.49 \pm 4.71 \text{ g kg}^{-1}$), a lower concentration was found in Březnice ($16.20 \pm 0.41 \text{ g kg}^{-1}$), and the lowest concentration was again recorded in Chmelná ($12.44 \pm 0.48 \text{ g kg}^{-1}$). However, the only significant difference was found between Březnice and Chmelná due to large concentration fluctuations in Běloky and Horní Bradlo. Phenolics concentrations in *Scirpus*




Fig. 3. Concentrations of total phenolic compounds in aboveground biomass of *Phalaris arundinacea* sampled during the period June–December 2016; data presented are means \pm SD

DW = dry biomass weight; a–c, A–C different letters indicate a significant difference at $\alpha = 0.05$ between the means




Fig. 4. Concentrations of total phenolic compounds in aboveground biomass of *Typha latifolia* sampled during the period June–December 2016; data presented are means \pm SD

DW = dry biomass weight; no statistical difference was found between the means

sylvaticus (Fig. 5E) were very high and varied between $31.30 \pm 2.24 \text{ g kg}^{-1}$ at Běloky and $21.31 \pm 0.07 \text{ g kg}^{-1}$ at Moraveč.

The concentrations of phenolics in *Glyceria maxima* (Fig. 5F) were low and varied only between $16.37 \pm 0.22 \text{ g kg}^{-1}$ at Louňovák and $12.02 \pm 0.03 \text{ g kg}^{-1}$ at Pařez (Fig. 5F). For *Juncus effusus*, no significant differences between four different localities were observed and values of phenolic compounds ranged from

$22.23 \pm 1.33 \text{ g kg}^{-1}$ in Běloky to $17.76 \pm 0.13 \text{ g kg}^{-1}$ in Moraveč (Fig. 5G).

DISCUSSION

There is a limited amount of information on phenolics content in wetland herbaceous macrophytes but the available data indicate that the concentrations are




Fig. 5. Average concentrations of total phenolic compounds in above-ground biomass of studied macrophytes harvested from various sites

DW = dry biomass weight; ^{a-c}different letters indicate a significant difference at $\alpha = 0.05$ between the means

lower as compared to wetland shrubs. The concentration found in *C. nigra* was very similar to the concentration of 31.4 g kg⁻¹ found in the same species in the Netherlands (Hoorens et al., 2003). Dorrepal (2005) reported phenolics concentrations between 7.3 and 19.8 g kg⁻¹ for five *Carex* species (*rotundata*, *vaginata*, *lasiocarpa*, *rostrate*, *acutiformis*) in Dutch fens indicating variability within the *Carex* genus. On the other hand, shrubs and scrubs usually exhibit higher concentrations of phenolics. For example, blueberry leaves are very rich source of phenolics. Routray, Orsat (2014) recorded phenolics concentration of 156 g kg⁻¹ in North American highbush blueberry (*Vaccinium corymbosum*) leaves. Also Dorrepal (2005) reported high phenolics concentrations in shrubs such as *Arctostaphylon alpinus* (black bearberry) – 301.6 g kg⁻¹, *Vaccinium uliginosum* (bog bilberry) – 118.5 g kg⁻¹ or *Vaccinium vitis-idaea* (lingonberry or cowberry) – 53 g kg⁻¹ sampled in Dutch fens. On the other hand, *Sphagnum* species have low phenolics concentrations (< 10 g kg⁻¹) and herbaceous wetland macrophytes seldom exceeded the concentration of 30 g kg⁻¹ (Dorrepal, 2005).

The total average value for all seven species found in our study was 18.59 ± 1.20 g kg⁻¹ DW (ranging from 9.02 to 28.39 g kg⁻¹ DW). Harrison et al. (2017) reported a very similar range of phenolics in 19 emergent wetland plants from 0.0 to 27.6 g kg⁻¹ DW with the average value 10.4 ± 6.20 g kg⁻¹ DW. Rejmankova (2016) recorded a relatively higher concentration of phenolics in tropical grown *Typha domingensis* leaves (phenolics: 22–46 g kg⁻¹; lignin: 19–38 g kg⁻¹) and relatively similar results in *Eleocharis cellulosa* stems (phenolics: 7–23 g kg⁻¹; lignin: 12–28 g kg⁻¹). In the same study it has also been shown that the difference in soluble phenolics concentrations of the greenhouse grown plants could be even higher – 87–133 g kg⁻¹ for *Typha* and 16–53 g kg⁻¹ for *Eleocharis*.

Our results also revealed that phenolics concentration varies between leaves and stems (Fig. 1). Hermann (1988) pointed out that the distribution of phenolics in the plant can be affected by light and some groups of phenolics can be mainly concentrated in tissues in parts exposed to more light. It has also been reported that various groups of phenolics are distributed differently in plant parts. Bujor et al. (2016) compared the diversity of phenolic compounds in leaves, stems, and fruits of bilberry (*Vaccinium myrtillus* L.). Although the variation of the total phenolic content was not so significant, there were large differences in the qualitative composition. Analyses showed the predominance of anthocyanins in fruits, caffeic acid derivatives in leaves, whereas flavonol oligomers represented more than half of the phenolic compounds in stems. In all bilberry parts, 106 phenolic compounds were evaluated. Of these 106 compounds, 62 were found in leaf (17 only in

leaves), 73 in stem (32 only in stems) and 40 in fruit extracts (9 only in fruits).

A variation of phenolics concentration depending on the harvesting time was also described by several authors for different plant species. Routray, Orsat (2014) evaluated the total phenolic content in *Vaccinium corymbosum* leaves harvested in May, July, September, and October. They observed the highest concentration in October and the lowest in July. The seasonal dynamics of polyphenols in submerged macrophyte *Myriophyllum verticillatum* L. (whorled water-milfoil) was evaluated by Bauer et al. (2009) during the growth seasons in four successive years. Total phenolic compounds contents significantly differed in the growing season and also even between years. Harrison et al. (2017) observed a significant decline of the foliar phenolic content in *Lythrum salicaria* (purple loosestrife) even during a very short period of sampling (from mid-July to the beginning of August).

The results supported the information that the concentrations of total phenolics in the aboveground biomass of herbaceous wetland macrophytes differ among species and localities as well. In addition, our study revealed that the phenolics concentrations vary between stems and leaves with concentrations being higher in leaves. Our study also showed that the concentration of phenolics decreases during the growing season. This is a quite important finding in relation to decomposition and potential carbon sequestration. Our further studies will focus on the relationship between the phenolic content in the aboveground biomass of wetland macrophytes and the nutrient status of the soil or sediment at the macrophyte stand. Also the relationship between the phenolics content and the decomposition rate will be evaluated. The future research will be aimed at the selection of plants that can be used for constructed wetlands treating effectively agricultural runoff and that can simultaneously contribute to carbon sequestration under the conditions of the Czech Republic.

CONCLUSION

During this study, concentrations of total phenolics were evaluated in seven common macrophytes in natural wetlands of the Czech Republic. The highest concentrations of phenolics were found in *Scirpus sylvaticus* and *Carex nigra*, while the lowest concentrations were observed in stems of *Phalaris arundinacea* and *Phragmites australis*. There was also a significant difference in phenolics concentrations in stems and leaves of these plants with leaves concentrations being higher. The evaluation of a seasonal dynamics of phenolics revealed that the concentration decreases throughout the year, however, the pattern varies among monitored species. The results also indicated a differ-

ence in the phenolics content in relation to sampling sites. The evaluation of the relationship between the nutrient status of soil and the phenolics in plants is the next phase of this research.

REFERENCES

- Aerts R, van Logtestijn RSP, Karlsson PS (2006): Nitrogen supply differentially affects litter decomposition rates and nitrogen dynamics of sub-arctic bog species. *Oecologia*, 146, 652–658. doi: 10.1007/s00442-005-0247-5.
- Barlocher F, Graca MAS (2005): Total phenolics. In: Graca MAS, Barlocher F, Gessner MO (eds): *Methods to study litter decomposition: a practical guide*. Springer, Dordrecht, 97–100. doi: 10.1007/1-4020-3466-0_14.
- Bauer N, Blaschke U, Beutler E, Gross EM, Jenett-Siems K, Siems K, Hilt S (2009): Seasonal and interannual dynamics of polyphenols in *Myriophyllum verticillatum* and their allelopathic activity on *Anabaena variabilis*. *Aquatic Botany*, 91, 110–116. doi: 10.1016/j.aquabot.2009.03.005.
- Boyd CE (1970): Losses of mineral nutrients during decomposition of *Typha latifolia*. *Archiv für Hydrobiologie*, 66, 511–517.
- Bravo L (1998): Polyphenols: chemistry, dietary sources, metabolism, and nutritional significance. *Nutrition Reviews*, 56, 317–333. doi: 10.1111/j.1753-4887.1998.tb01670.x.
- Bujor OC, Le Bourvellec C, Volf I, Popa VI, Dufour C (2016): Seasonal variations of the phenolic constituents in bilberry (*Vaccinium myrtillus* L.) leaves, stems and fruits, and their antioxidant activity. *Food Chemistry*, 213, 58–68. doi: 10.1016/j.foodchem.2016.06.042.
- Connor AM, Luby JJ, Tong CBS, Finn CE, Hancock JF (2002): Genotypic and environmental variation in antioxidant activity, total phenolic content, and anthocyanin content among blueberry cultivars. *Journal of the American Society for Horticultural Science*, 127, 89–97. doi: 10.17660/ActaHortic.2002.574.31.
- Dorrepaal E (2005): Plant growth-form and climate controls on production and decomposition in northern peatlands. Ph.D. Thesis, Vrije Universiteit Amsterdam.
- Gallagher JL (1978): Decomposition processes: summary and recommendation. In: Good RE, Whigham DF, Simpson RL (eds): *Freshwater wetlands. Ecological processes and management potential*. Academic Press, New York, 145–151.
- Giada MLR (2013): Food phenolic compounds: main classes, sources and their antioxidant power. In: Morales-Gonzalez JA (ed.): *Oxidative stress and chronic degenerative diseases – a role for antioxidants*. InTech, Rijeka.
- Harrison MM, Tyler AC, Hellquist CE, Pagano T (2017): Phenolic content of invasive and non-invasive emergent wetland plants. *Aquatic Botany*, 136, 146–154. doi: 10.1016/j.aquabot.2016.09.013.
- Hattenschwiler S, Vitousek PM (2000): The role of polyphenols in terrestrial ecosystem nutrient cycling. *Trends in Ecology and Evolution*, 15, 238–242. doi: 10.1016/S0169-5347(00)01861-9.
- Herrmann K (1988): On the occurrence of flavonol and flavone glycosides in vegetables. *Zeitschrift für Lebensmittel-Untersuchung und Forschung*, 186, 1–5. doi: 10.1007/BF01027170.
- Hoorens B, Aerts R, Stroetenga M (2003): Does initial litter chemistry explain litter mixture effects on decomposition? *Oecologia*, 137, 578–586. doi: 10.1007/s00442-003-1365-6.
- Horner JD, Gosz JR, Cates RG (1988): The role of carbon-based plant secondary metabolites in decomposition in terrestrial ecosystems. *The American Naturalist*, 132, 869–883. doi: 10.1086/284894.
- Howard LR, Clark JR, Brownmiller C (2003): Antioxidant capacity and phenolic content in blueberries as affected by genotype and growing season. *Journal of the Science of Food and Agriculture*, 83, 1238–1247. doi: 10.1002/jsfa.1532.
- Hutzler P, Fischbach R, Heller W, Jungblut TP, Reuber S, Schmitz R, Veit M, Weissenböck G, Schnitzler JP (1998): Tissue localization of phenolic compounds in plants by confocal laser scanning microscopy. *Journal of Experimental Botany*, 49, 953–965. doi: 10.1093/jxb/49.323.953.
- Jones CG, Hartley SE (1999): A protein competition model of phenolic allocation. *Oikos*, 86, 27–44. doi: 10.2307/3546567.
- Kala HK, Mehta R, Tandey R, Sen KK, Mandal V (2016): Ten years of research on phenolics (2005–2015): a status report. *Pacific Science Review A: Natural Science and Engineering*, 18, 1–4. doi: 10.1016/j.psra.2016.07.002.
- Kuiters AT (1990): Role of phenolic substances from decomposing forest litter in plant–soil interactions. *Acta Botanica Neerlandica*, 39, 329–348. doi: 10.1111/j.1438-8677.1990.tb01412.x.
- Lattanzio V (2013): Phenolic compounds: introduction. In: Ramawat KG, Mérillon JM (eds): *Natural products*. Springer-Verlag, Berlin, Heidelberg, 1543–1580.
- Murkovic M (2003): Phenolic compounds. In: Caballero B, Finglas P, Toldra F (eds): *Encyclopedia of Food Sciences and Nutrition*. Academic Press, San Diego, 4507–4514.
- Polunin NVC (1982): Processes in the decay of reed (*Phragmites australis*) litter in freshwater. In: Gopal B, Turner RE, Wetzel RG, Whigham DF (eds): *Wetlands: ecology and management*. International Scientific Publications, Jaipur, 293–304.
- Rejmankova E (2016): Phenolic content and growth of wetland macrophytes: is the allocation to secondary compounds driven by nutrient availability? *Folia Geobotanica*, 51, 239–250. doi: 10.1007/s12224-015-9227-7.
- Richardson CJ, Ferrell GM, Vaithianathan P (1999): Nutrient effects on stand structure, resorption efficiency, and secondary compounds in Everglades sawgrass. *Ecology*, 80, 2182–2192. doi: 10.2307/176902.
- Routray W, Orsat V (2014): Variation of phenolic profile and antioxidant activity of North American highbush blueberry

- leaves with variation of time of harvest and cultivar. *Industrial Crops and Products*, 62, 147–155. doi: 10.1016/j.indcrop.2014.08.020.
- Teal JM (1962): Energy flow in the salt marsh ecosystem in Georgia. *Ecology*, 43, 614–624. doi: 10.2307/1933451.
- Vermerris W, Nicholson R (2006a): The role of phenols in plant defense. In: Vermerris W, Nicholson R (eds): *Phenolic compound biochemistry*. Springer, Dordrecht, 211–234.
- Vermerris W, Nicholson R (2006b): Families of phenolic compounds and means of classification. In: Vermerris W, Nicholson R (eds): *Phenolic compound biochemistry*. Springer, Dordrecht, 1–34.
- Wang H, Richardson CJ, Ho M (2015): Dual controls on carbon loss during drought in peatlands. *Nature Climate Change*, 5, 584–587. doi: 10.1038/nclimate2643.
- Xu W, Shi L, Chan O, Li J, Casper P, Zou X (2013): Assessing the effect of litter species on the dynamic of bacterial and fungal communities during leaf decomposition in microcosm by molecular techniques. *PLoS ONE*, 8, e84613. doi: 10.1371/journal.pone.0084613.

Corresponding Author:

Prof. Ing. Jan Vymazal, CSc., Czech University of Life Sciences Prague, Faculty of Environmental Sciences, Kamýcká 129, 165 00 Prague 6-Suchbát, Czech Republic, phone: +420 224 383 825, e-mail: vymazal@yahoo.com
