



QUANTIFICATION OF FIRMICUTES, ACTINOBACTERIA, AND GAMMAPROTEOBACTERIA FROM BOHEMIAN HONEY*

Z. Hroncová^{1,2}, K. Konopásková¹, T. Volštatová¹, J. Killer^{1,3}

¹*Czech University of Life Sciences Prague, Faculty of Agrobiological Sciences, Department of Microbiology, Nutrition and Dietetics, Prague, Czech Republic*

²*Institute of Animal Science, Department of Genetics and Breeding of Farm Animals, Prague-Uhřetěves, Czech Republic*

³*Academy of Sciences of the Czech Republic, Institute of Animal Physiology and Genetics, Prague, Czech Republic*

Honey, which has been used as an ancient remedy for infected wounds, has been shown in laboratory studies to have antimicrobial action against a spectrum of bacteria and fungi. Because very little quantitative information exists on the microbiota of honey, the aim of this study was to quantify the Actinobacteria, Firmicutes, and Gammaproteobacteria groups in samples of honeydew honey and blossom honey from six regions in the Czech Republic, using quantitative real-time PCR analysis with specific primers based on the *16S rRNA* gene. Gammaproteobacteria and Firmicutes were clearly the most abundant, predominating Actinobacteria in both types of honey. Most of the Firmicutes were detected in samples from South Bohemia (mean gene copies per 1 g honey: 5.6×10^5) and Ústí nad Labem Region (3.7×10^5), which contained the lowest number of Gammaproteobacteria (15.5×10^3). The Actinobacteria were prevalent in samples from Plzeň (4.3×10^3) and Central Bohemia (5.4×10^3), where conversely the Firmicutes were least abundant. Honey thus contains bacterial species with probiotic activity and oligosaccharides which can act as prebiotics, suggesting that its incorporation into the human diet may potentially impart significant health benefits to consumers compared with ‘empty calories’ consumed as refined sugar.

Blossom, honeydew, Czech Republic, qPCR, microbiota



doi: 10.2478/sab-2018-0025

Received for publication on September 25, 2017

Accepted for publication on January 23, 2018

INTRODUCTION

Honey is the natural sweet substance produced by honey bees from the nectar of plants or from secretions of living parts of plants or excretions of plant-sucking insects on the living parts of plants, which the bees collect, transform by combining with specific substances of their own, and then deposit, dehydrate, store, and leave in the honey comb to ripen and mature (FAO, 2001). This sweet substance is most commonly consumed in its unpreserved state; that is, liquid, crystallized, or in the comb. In these forms, it is used as medicine, eaten as food, or incorporated as an ingredient into various food recipes. In confectionery production, honey is

still included in many traditional products which are consumed locally in considerable quantities and also exported. Honey is appreciated not only for its taste and flavour, but also for its high nutritive value and contribution to human health. There are many published reports describing a wide range of therapeutic effects (e.g. antibacterial, antiviral, antioxidative, and anti-inflammatory properties) of honey which are useful in stimulating the healing of wounds and burns and treating gastric ulcers and gastritis (Radwan et al., 1984; Jeddari et al., 1985; Subrahmanyam, 1991; Schramm et al., 2003; Al-Waili, 2004; Simon et al., 2006; van den Berg et al., 2008; Mandal, Mandal, 2011; Anthimidou, Mossialos,

* Supported by the National Agency for Agricultural Research - NAZV of the Ministry of Agriculture of Czech Republic, Project No. QJ1610248 and the Internal Grant Agency (CIGA) of the Czech University of Life Sciences Prague, Project No. 20162015.

2013; Tomblin et al., 2014; Almasaudi et al., 2015, 2017; Boyanova et al., 2015; Tsang et al., 2015). The high osmolality, acidity, and presence of hydrogen peroxide (H₂O₂) in natural honey make the growth of microorganisms in the substance difficult (Wahdan, 1998; Brudzynski, 2006; Kwakman et al., 2010; Mandal et al., 2010a, b; Aungzeb, Azim, 2011). Non-peroxide factors, such as lysozyme, phenolic acids, polyphenols, and flavonoids, also contribute to honey's antibacterial properties (Molan, 1992; Weston, 2000; Taormina et al., 2001; Brady et al., 2004; Kucuk et al., 2007; Hsieh et al., 2008; Al-Hindi et al., 2011; Al-Waili et al., 2011; Kwakman et al., 2011; Brudzynski et al., 2012; Kwakman, Zaat, 2012). In addition, because honey contains some propolis and bee pollen, part of its antimicrobial activity may be due to the presence of antimicrobial substances present in these components (Viuda-Martos et al., 2008; Redzic et al., 2011). Given these conditions, few microorganisms have the capacity to develop or remain in honey, and the microbes present are likely derived from primary or secondary sources of contamination. The primary sources of microbial contamination include pollen, the digestive tracts of honey bees, dirt, dust, air, and flowers. Secondary sources of microbial contamination in honey are humans, equipment, containers, wind, dust, etc. (Snowdon, Cliver, 1996; Olaitan et al., 2007). There are two recognised types of honey that differ in composition and which therefore should have different properties and bacterial spectra. Blossom honey (BH) is derived from the nectar of plants, whereas honeydew honey (HH) is derived mainly from the excretions of plant-sucking insects (Hemiptera) on the living parts of plants or the secretions of living parts of plants (FAO, 2001). Consumers have different demands for BH and HH; in many countries, nectar honey is valued more highly than HH, but in some countries, including the Czech Republic, HH is preferred (Sanz et al., 2005; Sanova et al., 2017). However, in spite of its usefulness, honey is known to contain certain microbes and is in fact described as a reservoir for microorganisms. The microbes present in honey are those that can withstand its concentrated sugar, acidity, and other antimicrobial components. Conventional microbiology and PCR-based studies have reported several species of cultivable and non-cultivable bacteria, yeasts, and filamentous fungi from honey, the compositions of which are tightly associated with the honey's botanical and geographical origins (Kropf et al., 2009; Castro-Vazquez et al., 2010; Kaskoniene, Venskutonis, 2010; Olivieri et al., 2012; Lazarevic et al., 2013; Sinacori et al., 2014). One of the widely used PCR techniques is quantitative PCR which analyses environmental and clinical microbiological samples. However, this technique has limitations in that it tends to underestimate or overestimate microbial counts, because its

main limitation is its inability to discriminate between dead and live cells including DNA of some of them that can be found in the environment (Masters et al., 1994; Wolffs et al., 2005; Sontakke et al., 2009; Maciel et al., 2011; Pathak et al., 2012; Li et al., 2013).

Despite the recent literature on the microbiota of honey, very little and only inconsistent information exists on the quantity of bacteria contained within this sweet substance. The aim of this study was therefore to compare samples of honeydew and blossom honey from six regions in the Czech Republic, using the quantitative real time polymerase chain reaction (qPCR) to quantify the Actinobacteria, Firmicutes, and Gammaproteobacteria groups specifically. Bacteria within these three groups are probiotic species or species with potential probiotic activity, which contaminate honey via the digestive tract of honey bees, nectar of flowers, or honeydew.

MATERIAL AND METHODS

Honey samples

The two honey types (22 samples of BH and 6 of HH) were sampled from six regions (Pardubice, Ústí nad Labem, Hradec Králové, Plzeň, South Bohemia, and Central Bohemia) in the Czech Republic (Table 1, Fig. 1). The samples and bee management represented traditional beekeeping practices in the Czech Republic. The honey samples were collected into disposable tubes and immediately frozen on dry ice, because freshness is related to storage and it is the problem of crystallized honey. Very few bacteria or microorganisms can survive in an environment like that, they




Fig. 1. Origin of the honey samples

Star indicates the place where blossom honey was sampled, a dot indicates the origin of honeydew honey, and a cross with a white dot indicates the place where both blossom honey and honeydew honey were sampled. Městečko u Křivoklátu (Central Bohemia) and Jaroměř (Hradec Králové) show only one cross because the two samples of blossom honey were from different beekeepers

Table 1. Places of origin and types of sampled honeys

| Region | Type of honey | Region | Type of honey |
|------------------------|---------------|-----------------------|---------------|
| Central Bohemia | | South Bohemia | |
| Hrabří | honeydew | Prachatice | honeydew |
| Hrabří | blossom | Dobrá Voda | honeydew |
| Městečko u Křivoklátu | blossom | Čimelice U Písku | blossom |
| Městečko u Křivoklátu | blossom | Hradec Králové | |
| Městečko u Křivoklátu | honeydew | Dvůr Králové | blossom |
| Řeřichy | blossom | Jaroměř | blossom |
| Senomaty | blossom | Jaroměř | blossom |
| Tmaň u Berouna | blossom | Přibyslav | blossom |
| Ústí nad Labem | | Přibyslav | honeydew |
| Kadaň | blossom | Pardubice | |
| Žatec | blossom | Břehy u Přelouče | blossom |
| Jirkov | blossom | Hlinsko | blossom |
| Plzeň | | Chvaletice | blossom |
| Klatovy | blossom | Pardubice | honeydew |
| Spůle | blossom | Rašovy u Přelouče | blossom |
| Horažďovice | blossom | Slatiňany | blossom |

just die, but qPCR detects also dead and live cell of bacteria. This means that freshness of honey does not influence the qPCR results. Every sample was weighed. Approximately 100–150 mg of the honey was used for individual isolation of the total bacterial

DNA, using the ZR Faecal DNA MiniPrep Kit (Zymo Research, Irvine, USA).

Quantitative real-time PCR analysis

Bacterial DNA was quantified using a MX3005P thermocycler (Stratagene, La Jolla, USA), based on the *16S rRNA* gene copy numbers with specific primers for Gammaproteobacteria (1080 γ F, γ 1202R), Firmicutes (928F-Firm, 1040FirmR), and Actinobacteria (Act920F3, Act1200R) (De Gregoris et al., 2011) (Table 2). IBM SPSS Statistics Version 20 (IBM, Armonk, USA) was used for the descriptive data analysis and for visualisation of the qPCR data.

RESULTS

The aim of this study was to quantify the probiotic species-containing Actinobacteria and Firmicutes as well as Gammaproteobacteria, as bacteria with potential probiotic activity against honey bees. However, the composition of honey changes depending on its botanical origin (HH or BH) and geographical region, because soil and climate characteristics determine the melliferous flora (Anklam, 1998; Rashid, Soltan, 2004; Castro-Vazquez et al., 2010). Therefore, we compared samples of HH and BH from six regions in the Czech Republic, using qPCR to quantify the three selected bacterial groups. Our results indicated that Gammaproteobacteria predominated in all the samples of honey (56.9% in HH vs 75.7% in




Figure 2. Quantification of bacterial DNA of Actinobacteria, Firmicutes, and Gammaproteobacteria

The number of copies of the *16S rRNA* gene per 1 g honey scaled to 100% is shown on the Y-axis

Table 2. Taxon-specific primer pairs used (De Gregoris et al., 2011)

| Target group | Primer | Sequence |
|---------------------|--------------------|--|
| Gammaproteobacteria | 1080γFγ1202R | TCGTCAGCTCGTGTGTGACGTAAGGGCCATGATG |
| Firmicutes | 928F-Firm1040FirmR | TGAAACTYAAAGGAATTGACGACCATGCACCACCTGTC |
| Actinobacteria | Act920F3Act1200R | TACGGCCGCAAGGCTATCRTCCCCACCTTCCTCCG |

BH), followed by Firmicutes (42.2% in HH vs 23.8% in BH) and Actinobacteria (0.9% in HH vs 0.6% in BH) in order of decreasing abundance (Fig. 2).

The HH samples contained more Firmicutes (mean gene copies per 1 g honey: 2.6×10^5) than the BH samples (1.2×10^5) (Fig. 3A). We also quantified the species of this bacterial phylum in samples from different Bohemian regions (Fig. 3B), with most of them being detected in samples from South Bohemia (5.6×10^5) and Ústí nad Labem (3.7×10^5). The least was in the honey from the region of Plzeň (2.1×10^4). As mentioned above, the most abundant group of bacteria was Gammaproteobacteria, likely contaminating the honey through the process of honey production, and it was higher in the BH samples (mean gene copies per 1 g honey: 3.9×10^5 vs 3.5×10^5 in HH samples) (Fig. 4A). However, this difference was not significant because the main source of Gammaproteobacteria is the digestive tract of honey bees. Samples of honey

from the region of Ústí nad Labem contained only 15.5×10^3 gene copies of Gammaproteobacteria (Fig. 4B). The differences in Gammaproteobacteria among the other five regions were not so considerable in comparison with those seen for Firmicutes and Actinobacteria. The least abundant group was Actinobacteria, which was more prevalent in HH (mean gene copies per 1 g honey: 5.8×10^3) than in BH (3.0×10^3) (Fig. 5A). The Actinobacteria were prevalent in samples from the regions of Plzeň (4.3×10^3) and Central Bohemia (5.4×10^3), where conversely the Firmicutes were the least abundant. Counts of Actinobacteria (5.6×10^2) were the lowest in South Bohemia (Fig. 5B).

DISCUSSION

Phylum Firmicutes, which includes two clusters (Firm4 and Firm5) that are largely restricted to the bee




Figure 3. Quantification of bacterial DNA of Firmicutes

(A) Quantitative determination (copies of the *16S rRNA* gene per 1 g honey) of Firmicutes in blossom honey and honeydew honey; values are means \pm SE. (B) Boxplot of quantitative real-time PCR (qPCR) data of the Firmicutes abundance in honey samples from six selected regions of the Czech Republic. The Y-axis shows copies of the *16S rRNA* gene per 1 g honey. Boxes show pooled data from honey samples from each region. The code of outlier 14 refers to the sample of honeydew honey from Pardubice, Region Pardubice

gut (Moran et al., 2012), was more prevalent in the samples of HH, which is generally characterised by higher values of electric conductivity, pH, and acidity (Terrab et al., 2003; Diez et al., 2004; Marini et al., 2004; Conti et al., 2007; Ouchemouk et

al., 2007; Manzanares et al., 2011). These conditions can create a suitable environment for lactobacilli. Moreover, the higher ash and oligosaccharide contents indicate potential prebiotic activity, increasing the populations of bifidobacteria and lactobacilli (Sanz




Figure 4. Quantification of bacterial DNA of Gammaproteobacteria

(A) Quantitative determination (copies of the 16S rRNA gene per 1 g honey) of Gammaproteobacteria in blossom honey and honeydew honey; values are means ± SE. (B) Boxplot of quantitative real-time PCR (qPCR) data of the Gammaproteobacteria abundance in honey samples from six selected regions of the Czech Republic. The Y-axis shows copies of the 16S rRNA gene per 1 g honey. Boxes show pooled data from samples of honey from each region




Figure 5. Quantification of bacterial DNA of Actinobacteria

(A) Quantitative determination (copies of the 16S rRNA gene per 1 g honey) of Actinobacteria in blossom honey and honeydew honey; values are means ± SE. (B) Boxplot of quantitative real-time PCR (qPCR) data of the Actinobacteria abundance in samples of honey from six selected regions of the Czech Republic. The Y-axis shows copies of the 16S rRNA gene per 1 g honey. Boxes show pooled data from samples of honey from each region. The code of outlier 7 refers to the sample of blossom honey from Přebyslav, Region Hradec Králové

et al., 2005). These can be beneficial not only as probiotics in honey, but also for the human gut microbiota, imparting nourishment benefits, such as fermentation ability and the break-down of nutrients to facilitate absorption of short-chain fatty acids, ions, amino acids, and vitamins; protective effects, preventing the invasion of pathogenic microorganisms; and trophic effects in the gut epithelium and digestive system (Anadon et al., 2016).

Although the BH samples contained more Gammaproteobacteria (Fig. 4A), this difference is not significant since the main source of these bacteria is the digestive tract of honey bees, and the honey is therefore contaminated through the process of honey production, when the honey bees ingest nectar and convert it with the help of enzymes. Besides these enzymes, some symbiotic microorganisms associated with the unique bee gut microbiota can also be incorporated into the honey. These consist mainly of eight bacterial phylogenotypes: two from the Alphaproteobacteria, two from the Gammaproteobacteria, two from *Lactobacillus*, one from *Bifidobacterium*, and one from the Betaproteobacteria (Martinson et al., 2011; Moran et al., 2012). Adult honey bees producing honey were shown to contain 5.1×10^7 gene copies of Gammaproteobacteria per 1 g of total digestive tract content (Hroncova et al., 2015). *Gilliamella apicola* and *Frischella perrara* are the most commonly occurring Gammaproteobacteria in the digestive tract of honey bees. Moreover, aphids producing honeydew were also shown to contain some symbiotic species of Gammaproteobacteria; namely, pea aphid secondary symbiont (PASS), pea aphid U-type symbiont (PAUS), pea aphid T-type symbiont (PABS) and *Buchnera* sp. (Unterman et al., 1989; Chen et al., 1996, 2000; Chen, Purcell, 1997; Fukatsu et al., 2000; Darby et al., 2001; Sandstrom et al., 2001; Tsuchida et al., 2002). However, our BH samples contained an excess of $\sim 3.9 \times 10^4$ gene copies of Gammaproteobacteria per 1 g of honey over that of the HH samples.

The least abundant bacterial group was Actinobacteria within the bifidobacteria cluster closely related to the honey bee gut (Rada et al., 1997; Jeyaprakash et al., 2003; Olofsson, Vasquez, 2008; Vasquez, Olofsson, 2009; Martinson et al., 2011; Moran et al., 2012). In general, these bacteria grow on rich media, consistent with their host-associated lifestyle, and require anaerobic or microaerophilic conditions, which is consistent with the likely lowered oxygen availability within the honey bee gut lumen compared with the conditions of the flower nectar. This caused the predominance of Actinobacteria in HH rather than in BH. However, HH has higher antioxidative and antibacterial properties (Prodoliet, Hischenhuber, 1998) which could be caused by the action of probiotic bacteria (bifidobacteria and lactobacilli), including the production of H_2O_2 , organic acids, bacteriocins, and strain-specific metabolites (Servin, 2004).

HH has an appreciably higher oligosaccharide content (Doner, 1977; Prodoliet, Hischenhuber, 1998), implying its potential prebiotic activity for increasing the populations of probiotic microbiota in the human gut (Sanz et al., 2005). Thus, daily intake of these new symbionts would be necessary to be able to populate the human body and maintain their benefits (Anadon et al., 2016).

CONCLUSION

Our results showed that HH contains more Firmicutes and Actinobacteria (both groups that contain beneficial bacteria) than BH which is rich in Gammaproteobacteria. The Actinobacteria and Gammaproteobacteria were the most abundant microbes in samples from the region of Central Bohemia. Conversely, the Firmicutes prevailed in honey from the region of South Bohemia. Although honey contains fewer microorganisms than other neutral foods, honeybee products nevertheless contain several lactic acid bacteria and bifidobacteria that act as beneficial probiotics when ingested, suggesting that incorporation of honey into the human diet or as a food ingredient may potentially impart significant health benefits to consumers. Therefore, our recommendation is to serve honey as a substitution for some of the 'empty calories' being consumed as refined sugar.

REFERENCES

- Al-Hindi RR, Bin-Masalam MS, El-Shahawi MS (2011): Antioxidant and antibacterial characteristics of phenolic extracts of locally produced honey in Saudi Arabia. *International Journal of Food Sciences and Nutrition*, 62, 513–517. doi: 10.3109/09637486.2010.550276.
- Almasaudi SB, Abbas AT, Al-Hindi RR, El-Shitany NA, Abdel-Dayem UA, Ali SS, Saleh RM, Al Jaouni SK, Kamal MA, Harakeh SM (2017): Manuka honey exerts antioxidant and anti-inflammatory activities that promote healing of acetic acid-induced gastric ulcer in rats. *Evidence-Based Complementary and Alternative Medicine*, 2017, 1–12. doi: 10.1155/2017/5413917.
- Almasaudi SB, El-Shitany NA, Abbas AT, Abdel-Dayem UA, Ali SS, Al Jaouni SK, Harakeh S (2015): Antioxidant, anti-inflammatory, and antiulcer potential of manuka honey against gastric ulcer in rats. *Oxidative Medicine and Cellular Longevity*, 2016, 1–10. doi: 10.1155/2016/3643824.
- Al-Waili NS (2004): Topical honey application vs. acyclovir for the treatment of recurrent herpes simplex lesions. *Medical Science Monitor*, 10, MT94–MT98.
- Al-Waili NS, Salom K, Butler G, Al Ghamdi AA (2011): Honey and microbial infections: a review supporting the use of

- honey for microbial control. *Journal of Medicinal Food*, 14, 1079–1096. doi: 10.1089/jmf.2010.0161.
- Anadon A, Martinez-Larranaga MR, Ares I, Martinez MA (2016): Prebiotics and probiotics: an assessment of their safety and health benefits. In: Watson RR, Preedy VR (ed): *Probiotics, prebiotics, and synbiotics. Bioactive foods in health promotion*. Elsevier, San Diego, USA, 3–23.
- Anklam E (1998): A review of the analytical methods to determine the geographical and botanical origin of honey. *Food Chemistry*, 63, 549–562. doi: 10.1016/S0308-8146(98)00057-0.
- Anthimidou E, Mossialos D (2013): Antibacterial activity of Greek and Cypriot honeys against *Staphylococcus aureus* and *Pseudomonas aeruginosa* in comparison to manuka honey. *Journal of Medicinal Food*, 16, 42–47. doi: 10.1089/jmf.2012.0042.
- Aurongzeb M, Azim MK (2011): Antimicrobial properties of natural honey: a review of literature. *Pakistan Journal of Biochemistry and Molecular Biology*, 44, 118–124.
- Boyanova L, Ilieva J, Gergova G, Vladimirov B, Nikolov R, Mitov I (2015): Honey and green/black tea consumption may reduce the risk of *Helicobacter pylori* infection. *Diagnostic Microbiology and Infectious Disease*, 82, 85–86. doi: 10.1016/j.diagmicrobio.2015.03.001.
- Brady N, Molan P, Bang L (2004): A survey of non-manuka New Zealand honeys for antibacterial and antifungal activities. *Journal of Apicultural Research*, 43, 47–52. doi: 10.1080/00218839.2004.11101109.
- Brudzynski K (2006): Effect of hydrogen peroxide on antibacterial activities of Canadian honeys. *Canadian Journal of Microbiology*, 52, 1228–1237. doi: 10.1139/w06-086.
- Brudzynski K, Abubaker K, Miotto D (2012): Unraveling a mechanism of honey antibacterial action: polyphenol/H₂O₂-induced oxidative effect on bacterial cell growth and on DNA degradation. *Food Chemistry*, 133, 329–336. doi: 10.1016/j.foodchem.2012.01.035.
- Castro-Vazquez L, Diaz-Maroto MC, De Torres C, Perez-Coello MS (2010): Effect of geographical origin on the chemical and sensory characteristics of chestnut honeys. *Food Research International*, 43, 2335–2340. doi: 10.1016/j.foodres.2010.07.007.
- Chen DQ, Purcell AH (1997): Occurrence and transmission of facultative endosymbionts in aphids. *Current Microbiology*, 34, 220–225. doi: 10.1007/s002849900172.
- Chen DQ, Campbell BC, Purcell AH (1996): A new Rickettsia from a herbivorous insect, the pea aphid *Acyrtosiphon pisum* (Harris). *Current Microbiology*, 33, 123–128. doi: 10.1007/s002849900086.
- Chen DQ, Montllor CB, Purcell AH (2000): Fitness effects of two facultative endosymbiotic bacteria on the pea aphid, *Acyrtosiphon pisum*, and the blue alfalfa aphid, *A. kondoi*. *Entomologia Experimentalis et Applicata*, 95, 315–323. doi: 10.1046/j.1570-7458.2000.00670.x.
- FAO (2001): Food and Agriculture Organization of the United Nations. Revised codex standard for honey (No. CODEX STAN 12–1981). www.fao.org/input/download/standards/310/cxs_012e.pdf. Accessed 13 July, 2018
- Conti ME, Stripeikis J, Campanella L, Cucina D, Tudino MB (2007): Characterization of Italian honeys (Marche Region) on the basis of their mineral content and some typical quality parameters. *Chemistry Central Journal*, 1, 14. doi: 10.1186/1752-153X-1-14.
- Darby AC, Birkle LM, Turner SL, Douglas AE (2001): An aphid-borne bacterium allied to the secondary symbionts of whitefly. *FEMS Microbiology Ecology*, 36, 43–50. doi: 10.1111/j.1574-6941.2001.tb00824.x.
- De Gregoris TB, Aldred N, Clare AS, Burgess JG (2011): Improvement of phylum- and class-specific primers for real-time PCR quantification of bacterial taxa. *Journal of Microbiological Methods*, 86, 351–356. doi: 10.1016/j.mimet.2011.06.010.
- Diez MJ, Andres C, Terrab A (2004): Physicochemical parameters and pollen analysis of Moroccan honeydew honeys. *International Journal of Food Science and Technology*, 39, 167–176. doi: 10.1046/j.0950-5423.2003.00769.x.
- Doner LW (1977): The sugars of honey – a review. *Journal of the Science of Food and Agriculture*, 28, 443–456. doi: 10.1002/jsfa.2740280508.
- Fukatsu T, Nikoh N, Kawai R, Koga R (2000): The secondary endosymbiotic bacterium of the pea aphid *Acyrtosiphon pisum* (Insecta: Homoptera). *Applied and Environmental Microbiology*, 66, 2748–2758. doi: 10.1128/AEM.66.7.2748-2758.2000.
- Hroncova Z, Havlik J, Killer J, Doskocil I, Tyl J, Kamler M, Titera D, Hakl J, Mrazek J, Bunesova V (2015): Variation in honey bee gut microbial diversity affected by ontogenetic stage, age and geographic location. *PLoS ONE*, 10, e0118707. doi: 10.1371/journal.pone.0118707.
- Hsieh MC, Shen YJ, Kuo YH, Hwang LS (2008): Antioxidative activity and active components of longan (*Dimocarpus longan* Lour.) flower extracts. *Journal of Agricultural and Food Chemistry*, 56, 7010–7016. doi: 10.1021/jf801155j.
- Jeddar A, Kharsany A, Ramsaroop UG, Bhamjee A, Haffejee IE, Moosa A (1985): The antibacterial action of honey. An in vitro study. *South African Medical Journal*, 67, 257–258.
- Jeyaprakash A, Hoy MA, Allsopp MH (2003): Bacterial diversity in worker adults of *Apis mellifera capensis* and *Apis mellifera scutellata* (Insecta: Hymenoptera) assessed using 16S rRNA sequences. *Journal of Invertebrate Pathology*, 84, 96–103. doi: 10.1016/j.jip.2003.08.007.
- Kaskoniene V, Venskutonis PR (2010): Floral markers in honey of various botanical and geographic origins: a review. *Comprehensive Reviews in Food Science and Food Safety*, 9, 620–634. doi: 10.1111/j.1541-4337.2010.00130.x.
- Kropf U, Bertonecelj J, Korosec M, Necemer M, Kump P, Ogrinc N (2009): Geographical origin of Slovenian multifloral and forest honey. *Apiacta*, 44, 33–42.
- Kucuk M, Kolayli S, Karaoglu S, Ulusoy E, Baltaci C, Candan F (2007): Biological activities and chemical composition of

- three honeys of different types from Anatolia. *Food Chemistry*, 100, 526–534. doi: 10.1016/j.foodchem.2005.10.010.
- Kwakman PHS, Zaat SAJ (2012): Antibacterial components of honey. *IUBMB Life*, 64, 48–55. doi: 10.1002/iub.578.
- Kwakman PHS, Te Velde AA, De Boer L, Speijer D, Vandenbroucke-Grauls CMJE, Zaat SAJ (2010): How honey kills bacteria. *The FASEB Journal*, 24, 2576–2582. doi: 10.1096/fj.09-150789.
- Kwakman PHS, Te Velde AA, De Boer L, Vandenbroucke-Grauls CMJE, Zaat SAJ (2011): Two major medicinal honeys have different mechanisms of bactericidal activity. *PLoS ONE*, 6, e17709. doi: 10.1371/journal.pone.0017709.
- Lazarevic KB, Trifkovic JD, Andric FL, Tesic ZL, Anđelkovic IB, Radovic DI, Nedic NM, Milojkovic-Opšenica DM (2013): Quality parameters and pattern recognition methods as a tool in tracing regional origin of multifloral honey. *Journal of the Serbian Chemical Society*, 78, 1875–1892. doi: 10.2298/JSC130701099L.
- Li M, Zhao G, Liu J, Gao X, Zhang Q (2013): Effect of different heat treatments on the degradation of *Salmonella* nucleic acid. *Journal of Food Safety*, 33, 536–544. doi: 10.1111/jfs.12086.
- Maciel BM, Dias JC, Romano CC, Srianganathan N, Brendel M, Rezende RP (2011): Detection of *Salmonella* Enteritidis in asymptomatic carrier animals: comparison of quantitative real-time PCR and bacteriological culture methods. *Genetics and Molecular Research*, 10, 2578–2588. doi: 10.4238/2011.
- Mandal MD, Mandal S (2011): Honey: its medicinal property and antibacterial activity. *Asian Pacific Journal of Tropical Biomedicine*, 1, 154–160. doi: 10.1016/S2221-1691(11)60016-6.
- Mandal S, Debmandal M, Pal NK, Saha K (2010a): Antibacterial activity of honey against clinical isolates of *Escherichia coli*, *Pseudomonas aeruginosa* and *Salmonella enterica* serovar Typhi. *Asian Pacific Journal of Tropical Medicine*, 3, 961–964. doi: 10.1016/S1995-7645(11)60009-6.
- Mandal S, Debmandal M, Pal NK, Saha K (2010b): Synergistic anti-*Staphylococcus aureus* activity of amoxicillin in combination with *Emblica officinalis* and *Nymphae odorata* extracts. *Asian Pacific Journal of Tropical Medicine*, 3, 711–714. doi: 10.1016/S1995-7645(10)60171-X.
- Manzanares AB, Garcia ZH, Galdon BR, Rodriguez ER, Romero CD (2011): Differentiation of blossom and honeydew honeys using multivariate analysis on the physicochemical parameters and sugar composition. *Food Chemistry*, 126, 664–672. doi: 10.1016/j.foodchem.2010.11.003.
- Marini F, Magri AL, Balestrieri F, Fabretti F, Marini D (2004): Supervised pattern recognition applied to the discrimination of the floral origin of six types of Italian honey samples. *Analytica Chimica Acta*, 515, 117–125. doi: 10.1016/j.aca.2004.01.013.
- Martinson VG, Danforth BN, Minckley RL, Rueppell O, Tingek S, Moran NA (2011): A simple and distinctive microbiota associated with honey bees and bumble bees. *Molecular Ecology*, 20, 619–628. doi: 10.1111/j.1365-294X.2010.04959.x.
- Masters CI, Shallcross JA, Mackey BM (1994): Effect of stress treatments on the detection of *Listeria monocytogenes* and enterotoxigenic *Escherichia coli* by the polymerase chain reaction. *Journal of Applied Microbiology*, 77, 73–79. doi: 10.1111/j.1365-2672.1994.tb03047.x.
- Molan PC (1992): The antibacterial activity of honey: 2. Variation in the potency of the antibacterial activity. *Bee World*, 73, 59–76. doi: 10.1080/0005772X.1992.11099118.
- Moran NA, Hansen AK, Powell JE, Sabree ZL (2012): Distinctive gut microbiota of honey bees assessed using deep sampling from individual worker bees. *PLoS ONE*, 7, e36393. doi: 10.1371/journal.pone.0036393.
- Olaitan PB, Adeleke OE, Iyabo OO (2007): Honey: a reservoir for microorganisms and an inhibitory agent for microbes. *African Health Sciences*, 7, 159–165. doi: 10.5555/afhs.2007.7.3.159.
- Olivieri C, Marota I, Rollo F, Luciani S (2012): Tracking plant, fungal, and bacterial DNA in honey specimens. *Journal of Forensic Sciences*, 57, 222–227. doi: 10.1111/j.1556-4029.2011.01964.x.
- Olofsson TC, Vasquez A (2008): Detection and identification of a novel lactic acid bacterial flora within the honey stomach of the honeybee *Apis mellifera*. *Current Microbiology*, 57, 356–363. doi: 10.1007/s00284-008-9202-0.
- Ouchemoukh S, Louaileche H, Schweitzer P (2007): Physicochemical characteristics and pollen spectrum of some Algerian honeys. *Food Control*, 18, 52–58. doi: 10.1016/j.foodcont.2005.08.007.
- Pathak S, Awuh JA, Leversen NA, Flo TH, Asjo B (2012): Counting mycobacteria in infected human cells and mouse tissue: a comparison between qPCR and CFU. *PLoS ONE*, 7, e34931. doi: 10.1371/journal.pone.0034931.
- Prodoliet J, Hischenhuber C (1998): Food authentication by carbohydrate chromatography. *Zeitschrift für Lebensmitteluntersuchung und-Forschung A*, 207, 1–12. doi: 10.1007/s002170050286.
- Rada V, Machova M, Huk J, Marounek M, Duskova D (1997): Microflora in the honeybee digestive tract: counts, characteristics and sensitivity to veterinary drugs. *Apidologie*, 28, 357–365. doi: 10.1051/apido:19970603.
- Radwan SS, El-Essawy AA, Sarhan MM (1984): Experimental evidence for the occurrence in honey of specific substances active against microorganisms. *Zentralblatt für Mikrobiologie*, 139, 249–255. doi: 10.1016/S0232-4393(84)80047-5.
- Rashed MN, Soltan ME (2004): Major and trace elements in different types of Egyptian mono-floral and non-floral bee honeys. *Journal of Food Composition and Analysis*, 17, 725–735. doi: 10.1016/j.jfca.2003.10.004.
- Redzic S, Kurtagic H, Prazina N, Tuka M, Avdagic T (2011): The antimicrobial activity of honey in relation to the composition of pollen (Bosnia-Herzegovina, W. Balkan). *Planta Medica*, 77, SL62. doi: 10.1055/s-0031-1282185.

- Sandstrom JP, Russell JA, White JP, Moran NA (2001): Independent origins and horizontal transfer of bacterial symbionts of aphids. *Molecular Ecology*, 10, 217–228. doi: 10.1046/j.1365-294X.2001.01189.x.
- Sanova P, Svobodova J, Hrubcova B, Serakova P (2017): Segmentation of honey buyers' behaviour by conjoint analysis. *Scientia Agriculturae Bohemica*, 48, 55–62. doi: 10.1515/sab-2017-0008.
- Sanz ML, Gonzalez M, De Lorenzo C, Sanz J, Martinez-Castro I (2005): A contribution to the differentiation between nectar honey and honeydew honey. *Food Chemistry*, 91, 313–317. doi: 10.1016/j.foodchem.2004.06.013.
- Schramm DD, Karim M, Schrader HR, Holt RR, Cardetti M, Keen CL (2003): Honey with high levels of antioxidants can provide protection to healthy human subjects. *Journal of Agricultural and Food Chemistry*, 51, 1732–1735. doi: 10.1021/jf025928k.
- Servin AL (2004): Antagonistic activities of lactobacilli and bifidobacteria against microbial pathogens. *FEMS Microbiology Reviews*, 28, 405–440. doi: 10.1016/j.femsre.2004.01.003.
- Simon A, Sofka K, Wiszniewsky G, Blaser G, Bode U, Fleischhack G (2006): Wound care with antibacterial honey (Medihoney) in pediatric hematology–oncology. *Supportive Care in Cancer*, 14, 91–97. doi: 10.1007/s00520-005-0874-8.
- Sinacori M, Francesca N, Alfonzo A, Cruciata M, Sannino C, Settanni L, Moschetti G (2014): Cultivable microorganisms associated with honeys of different geographical and botanical origin. *Food Microbiology*, 38, 284–294. doi: 10.1016/j.fm.2013.07.013.
- Snowdon JA, Cliver DO (1996): Microorganisms in honey. *International Journal of Food Microbiology*, 31, 1–26. doi: 10.1016/0168-1605(96)00970-1.
- Sontakke S, Cadenas M, Maggi R, Diniz P, Breitschwerdt EB (2009): Use of broad range 16S rDNA PCR in clinical microbiology. *Journal of Microbiological Methods*, 76, 217–225. doi: 10.1016/j.mimet.2008.11.002.
- Subrahmanyam M (1991): Topical application of honey in treatment of burns. *British Journal of Surgery*, 78, 497–498. doi: 10.1002/bjs.1800780435.
- Taormina PJ, Niemira BA, Beuchat LR (2001): Inhibitory activity of honey against foodborne pathogens as influenced by the presence of hydrogen peroxide and level of antioxidant power. *International Journal of Food Microbiology*, 69, 217–225. doi: 10.1016/S0168-1605(01)00505-0.
- Terrab A, Gonzalez AG, Diez MJ, Heredia FJ (2003): Characterisation of Moroccan unifloral honeys using multivariate analysis. *European Food Research and Technology*, 218, 88–95. doi: 10.1007/s00217-003-0797-x.
- Tomblin V, Ferguson LR, Han DY, Murray P, Schlothauer R (2014): Potential pathway of anti-inflammatory effect by New Zealand honeys. *International Journal of General Medicine*, 7, 149–158. doi: 10.2147/IJGM.S45839.
- Tsang KK, Kwong EWY, Woo KY, To TSS, Chung JWY, Wong TKS (2015): The anti-inflammatory and antibacterial action of nanocrystalline silver and manuka honey on the molecular alternation of diabetic foot ulcer: a comprehensive literature review. *Evidence-Based Complementary and Alternative Medicine*, 2015, 1–19. doi: 10.1155/2015/218283.
- Tsuchida T, Koga R, Shibao H, Matsumoto T, Fukatsu T (2002): Diversity and geographic distribution of secondary endosymbiotic bacteria in natural populations of the pea aphid, *Acyrtosiphon pisum*. *Molecular Ecology*, 11, 2123–2135. doi: 10.1046/j.1365-294X.2002.01606.x.
- Unterman BM, Baumann P, McLean DL (1989): Pea aphid symbiont relationships established by analysis of 16S rRNAs. *Journal of Bacteriology*, 171, 2970–2974. doi: 10.1128/jb.171.6.2970-2974.1989.
- Van den Berg AJJ, Van den Worm E, Quarles van Ufford HC, Halkes SBA, Hoekstra MJ, Beukelman CJ (2008): An in vitro examination of the antioxidant and anti-inflammatory properties of buckwheat honey. *Journal of Wound Care*, 17, 172–178. doi: 10.12968/jowc.2008.17.4.28839.
- Vasquez A, Olofsson TC (2009): The lactic acid bacteria involved in the production of bee pollen and bee bread. *Journal of Apicultural Research*, 48, 189–195. doi: 10.3896/IBRA.1.48.3.07.
- Viuda-Martos M, Ruiz-Navajas Y, Fernandez-Lopez J, Perez-Alvarez JA (2008): Functional properties of honey, propolis, and royal jelly. *Journal of Food Science*, 73, 117–124. doi: 10.1111/j.1750-3841.2008.00966.x.
- Wahdan HAL (1998): Causes of the antimicrobial activity of honey. *Infection*, 26, 26–31. doi: 10.1007/BF02768748.
- Weston RJ (2000): The contribution of catalase and other natural products to the antibacterial activity of honey: a review. *Food Chemistry*, 71, 235–239. doi: 10.1016/S0308-8146(00)00162-X.
- Wolffs P, Norling B, Radstrom P (2005): Risk assessment of false-positive quantitative real-time PCR results in food, due to detection of DNA originating from dead cells. *Journal of Microbiological Methods*, 60, 315–323. doi: 10.1016/j.mimet.2004.10.003.

Corresponding Author:

Ing. Zuzana Hroncová, Ph.D., Czech University of Life Sciences Prague, Faculty of Agrobiology, Food and Natural Resources, Department of Microbiology, Nutrition and Dietetics, Kamýcká 129, 165 00 Prague 6-Suchbát, Czech Republic, phone: +420 224 382 669, e-mail: hroncovaz@af.czu.cz
